

HETEROPOLY ACID SOLUTION TYPE

NPM-40 (PHOSPHOMOLYBDIC ACID SOLUTION)	2
NPW-40 (PHOSPHOTUNGSTIC ACID SOLUTION)	3
NSW-40 (SILICOTUNGSTIC ACID SOLUTION)	4

NPM-40 (PHOSPHOMOLYBDIC ACID SOLUTION)



- **OUTLINE**

SPECIFIC GRAVITY (20 °C): approx. 1.45

MoO₃ CONTENT: approx. 37%

- **APPLICATIONS**

Acid catalyst, Oxidation catalyst

- **SPECIFICATIONS**

MoO ₃	(37±2)%
Na	0.01% Max.
Cl	0.002% Max.
SO ₄	0.005% Max.
Fe	0.003% Max.
Pb	0.004% Max.
NH ₄	0.005% Max.

NPW-40 (PHOSPHOTUNGSTIC ACID SOLUTION)



- **OUTLINE**

SPECIFIC GRAVITY (20 °C): approx. 1.55

WO₃ CONTENT: approx. 40%

- **APPLICATIONS**

Acid catalyst, Oxidation catalyst

- **SPECIFICATIONS**

WO ₃	(40±1)%
Na	0.01% Max.
Cl	0.002% Max.
SO ₄	0.005% Max.
Fe	0.003% Max.
Pb	0.004% Max.
NH ₄	0.005% Max.

NSW-40 (SILICOTUNGSTIC ACID SOLUTION)



- **OUTLINE**

SPECIFIC GRAVITY (20 °C): approx. 1.55

WO₃ CONTENT: approx. 40%

- **APPLICATIONS**

Acid catalyst

- **SPECIFICATIONS**

WO ₃	(40±2)%
Na	0.01% Max.
Cl	0.002% Max.
SO ₄	0.005% Max.
Fe	0.003% Max.
Pb	0.004% Max.
NH ₄	0.005% Max.



NIPPON INORGANIC COLOUR & CHEMICAL CO., LTD.

3-14-1 Funado, Itabashi-ku, Tokyo, 174-0041

(Main Office & Plant)

TEL : +81-3-3966-1277

FAX : +81-3-3967-9266

(Sales Div.)

TEL : +81-3-3966-2412

FAX : +81-3-3966-2680